



South Ruby Mountains, NV




Woodland Invasion (expansion, encroachment, reclamation)



1904



1998




Are we getting what we expect?

Short-term response by bird communities to treatments in pinyon-juniper woodlands


Steve Knick and Steve Hanser
USGS

Matthias Leu
College of William and Mary



USGS Disclaimer: Results presented are preliminary and have not been peer-reviewed

Space – Time Relationships



*So, Nat'ralists observe, a Flea
Hath smaller Fleas that on him prey,
And these have smaller Fleas to bit 'em,
And so proceed ad infinitum.*

Jonathan Swift 1733

Greater Sage-Grouse


The Problems

- Large home range
- Long lived
- Low reproductive rate



Terry Steele

Spatial scale – Disturbance/territory



Brewer's Sparrow

Black-throated Sparrow

Gray Flycatcher

Grasshopper Sparrow

Green-tailed Towhee

Horned Lark

Lark Sparrow

Loggerhead Shrike

Sage Sparrow

Sage Thrasher

Savannah Sparrow


Vesper Sparrow

Western Meadowlark




Photos: BNA Online


Spatial scale – Disturbance/territory




Spatial scale – Disturbance/territory



Spatial scale – Disturbance/territory







Objectives

- **Delineate the organization of bird communities**
- **Identify the environmental gradients underlying bird communities**
- **Use changes in bird community to interpret the effects of management treatment**



Bird Community




Study Sites and Woodland Ecosystems



Woodland Network Site Ordination



Sites and Environmental Gradients





Study Design

- Paired study sites
 - Treatment and Control
- Breeding bird point transects
- Vegetation plots
- Ordinations (CCA)



Point Transects

Castlehead Site, ID





Vegetation Sampling


- 2 30 x 30 m sampling plots per point count
- Vegetation ground cover





Ordination























courtesy Jeff Rose





Five Creeks



Five Creeks

- Burned in Fall 2008
 - Pre-treatment data collected in 2008
- 5300-6300 ft elevation
- Western juniper, mountain big sagebrush, and low sagebrush with a mixture of native grasses (Sandberg bluegrass, Idaho fescue, and squirrel tail).
- 83% of veg point associated with PC's burned
- 76.6% of area within burn plots was actually burned

Pre-treatment

2008

Post-treatment

2009

2010

2011



Five Creeks



2005 Pre-Treatment



2009 Post-Treatment

Five Creeks



2005 Pre-Treatment



2009 Post-Treatment


2008

Five Creeks


2011




Five Creeks




Five Creeks




Five Creeks




Five Creeks




Five Creeks





Five Creeks



Sagebrush obligates



Gray flycatcher




Five Creeks Bird Response


- Species richness remained the same

Winners*	Losers*	No Change
American Robin	Chipping Sparrow	Brewer's Sparrow
Dark-eyed Junco	Gray Flycatcher	Cassin's Finch
Northern Flicker	Green-tailed Towhee	Dusky Flycatcher
Vesper Sparrow		House Wren
		Mountain Bluebird
		Rock Wren
		White-crowned Sparrow
		Western Meadowlark

*P < 0.05




Onaqui



- Western Juniper
- Pinyon - Juniper
- Utah Juniper


Onaqui

- Burned in Fall 2006
 - Pre-treatment data in 2006
- 5500-6200 ft elevation
- Utah Juniper, Wyoming big sagebrush, black sagebrush, bluebunch wheatgrass, Sandberg bluegrass, Indian ricegrass, cheatgrass, and various native wildflowers
- 65% of veg points associated with PC's burned
- 44.7% of area within burn plots was actually burned



Onaqui

Approximate burn delineation



Onaqui




2006





2007

Onaqui




Onaqui







Onaqui




Onaqui




Onaqui




Onaqui




Onaqui




Onaqui





Onaqui



Sagebrush Obligates



Gray flycatcher




Onaqui

Bird Response


- Species richness remained the same


Winners*	Losers*	No Change
	Bewick's Wren	Brewer's Sparrow
	Gray Flycatcher	Blue-gray Gnatcatcher
	Lark Sparrow	Spotted Towhee
	Horned Lark	House Finch
		Chipping Sparrow
		Juniper Titmouse
		Black-throated Gray Warbler
		Sage Sparrow
		Sage Thrasher
		Western Meadowlark


Western meadowlark




Bird Community









Correlations – Axis 1

	BBS	Control
Control	0.69 $p < 0.01$	
Treatment	ns	ns

Conclusions

- Treatment and control differ in bird community structure and trajectory
- Short-term response to treatment may be primarily a function of habitat loss
- Habitat improvements for sagebrush obligates may be long-term
- Regional dynamics also are important

Acknowledgements

- Joint Fire Science Program
- SageSTEP project 
- US Geological Survey 
- Bureau of Land Management and US Forest Service for logistical support
- Field Technicians



courtesy Jeff Rose